

KOBE-FHD-95-03

April 5 1995

TWO-SPIN ASYMMETRY OF HEAVY QUARKONIA PRODUCTIONS AT RHIC ENERGY

T. YAMANISHI [†]

*Graduate School of Science and Technology,
Kobe University, Kobe, Japan 657*

and

T. MORII ^{††}, S. TANAKA

*Faculty of Human Development, Division of
Sciences for Natural Environment,
Kobe University, Kobe, Japan 657*

ABSTRACT

Hadroproductions of charmonium and bottomonium χ_J in polarized pp collisions at RHIC energies are studied. Two-spin asymmetries $A_{LL}^{\chi_J}(pp)$ for these reactions are very good parameters for examining the polarized gluon distributions in a proton.

1 Introduction

Relativistic Heavy Ion Collider (RHIC)[1] which is designed to have a beam polarization of about 70 % with a luminosity of $2 \times 10^{32} \text{ cm}^{-2} \text{ sec}^{-1}$ at an energy of $\sqrt{s} = 50 - 500 \text{ GeV}$ will give us fruitful informations on the spin structure of proton. Recent measurements of the spin-dependent proton structure function $g_1^p(x, Q^2)$ by EMC[2] and SMC Collaboration[3], which are far from the Ellis-Jaffe prediction[4], confirmed again the suggestion that very little of the proton spin is carried by quarks. Many ideas have been proposed so far to solve this problem. One of the interesting approach is that the gluon plays a crucial role on the nucleon spin structure[5]. But the magnitude ΔG and distribution $\delta G(x)$ of the gluon polarization inside a proton is not known yet. In order to improve our understanding of the nucleon structure, it is necessary to measure ΔG and $\delta G(x)$ in other reactions which are sensitive to the gluon polarization.

[†]Presented in the 1995 Lake Louise Winter Institute at Lake Louise, Canada on 19-25 February.
E-mail yamanisi@natura.h.kobe-u.ac.jp

^{††}E-mail morii@kobe-u.ac.jp

Among various reactions, the χ_J -productions in hadron-hadron collisions are interesting candidates. In this paper, we are concerned with the χ_J -hadroproductions at low- p_T regions. The cross sections of these processes are expected to depend sensitively on the shape of polarized gluon distributions because the gluon-gluon fusion is the dominant mechanism at the lowest order of QCD (see Figure 1). Although these productions have been carefully studied so far by several people in order to extract unpolarized gluon distributions $G(x)$ and test the perturbative QCD[6], we are interested in the behavior of the polarized gluon distributions $\delta G(x)$.

So far, various types of the polarized gluon distribution functions $\delta G(x)$ have been proposed: some of them have large ΔG ($\simeq 5 - 6$)[7, 8, 9, 10] and others have small ΔG ($\lesssim 2 - 3$)[8, 9, 11, 12], where $\Delta G = \int_0^1 \delta G(x) dx$. Here we take the following typical types of $x\delta G(x)$ to examine the effect on the χ_J -hadroproductions:

(a) MTY model [10] :

$$x\delta G(x, Q^2 = 10\text{GeV}^2) = 9.52x^{0.4}(1-x)^{17} , \quad (1)$$

with $\Delta G(Q_{SMC}^2) = 5.32$,

(b) Cheng-Lai model [8] :

$$x\delta G(x, Q^2 = 10\text{GeV}^2) = 3.34x^{0.31}(1-x)^{5.06}(1-0.177x) , \quad (2)$$

with $\Delta G(Q_{SMC}^2) = 5.64$,

(c) BBS model [11] :

$$x\delta G(x, Q^2 = 4\text{GeV}^2) = 0.281 \left\{ (1-x)^4 - (1-x)^6 \right\} \\ + 1.1739 \left\{ (1-x)^5 - (1-x)^7 \right\} , \quad (3)$$

with $\Delta G(Q_{SMC}^2) = 0.53$,

(d) no gluon polarization model :

$$x\delta G(x, Q^2 = 10\text{GeV}^2) = 0 , \text{ with } \Delta G(Q_{SMC}^2) = 0 , \quad (4)$$

where Q_{SMC}^2 is taken to be 10GeV^2 . Among these distributions, ΔG of types (a) and (b) are large while those of types (c) and (d) are small and zero, respectively. The x -dependence of $x\delta G(x, Q^2)$ at $Q^2 = 10\text{GeV}^2$ and $\delta G(x, Q^2)/G(x, Q^2)$ which are evolved up to $Q^2 = M_{\chi_2(c\bar{c})}^2$ and $M_{\chi_2(b\bar{b})}^2$ by the Altarelli-Parisi equations, is depicted in Figures 2(A), (B) and (C), respectively. Most people[7] who assume large ΔG take type (b) as the $x\delta G$. As shown in Figure 2(A), the $x\delta G(x)$ of type (b) has a peak

at $x \approx 0.05$ and gradually decreases with increasing x while that of (a) has a sharp peak at $x \approx 0.04$ and rapidly decreases with x . It is remarkable that the distribution of type (a) is consistent with the experimental data of not only two-spin asymmetries $A_{LL}^{\pi^0}(\bar{p}p)$ for the inclusive π^0 -production[10, 13] but also recent $A_{LL}^{\gamma\gamma}(pp)$ for the inclusive multi- γ pair production[14] measured by the E581/704 Collaboration at Fermilab by using longitudinally polarized proton (antiproton) beams and longitudinally polarized proton targets, even though it has large $\Delta G (= 5.32)$ in a proton. Here we examine above four types of polarized gluon distribution functions to discuss low- p_T χ_J -hadroproductions, though the type (b) has been already ruled out by the data on two-spin asymmetries[15].

2 Two-spin asymmetry A_{LL}

Let us start by defining an interesting parameter called a two-spin asymmetry for χ_J -hadroproductions $A_{LL}^{\chi_J}$, which is given by

$$\begin{aligned} A_{LL}^{\chi_J} &= \frac{[d\sigma_{++} - d\sigma_{+-} + d\sigma_{--} - d\sigma_{-+}]}{[d\sigma_{++} + d\sigma_{+-} + d\sigma_{--} + d\sigma_{-+}]} \\ &= \frac{d\Delta\sigma_{\chi_J}}{d\sigma_{\chi_J}}. \end{aligned} \quad (5)$$

Here $d\sigma_{+-}$, for instance, denotes that the helicity of one beam particle is positive and the one of the other is negative. Although several people have investigated $A_{LL}^{\chi_J}$ as a function of \sqrt{s} [16], the experimental data are available more easily as a function of the longitudinal fraction of produced particle x_L . Then we investigate how the x_L -dependence of $A_{LL}^{\chi_J}$ for χ_J -hadroproductions in polarized pp collisions are affected by polarized gluon distribution functions.

Since the dominant source is the gluon-gluon fusion as shown in Figure 1, the reaction is very sensitive to polarized gluon distributions in a proton. The χ_1 state cannot be produced by this reaction due to Yang's theorem. The spin-dependent and spin-independent subprocess cross sections $\Delta\hat{\sigma}_{\chi_J}$ and $\hat{\sigma}_{\chi_J}$ for the subprocess $G + G \rightarrow \chi_{0,2}$ are straightforwardly calculated as[17]

$$\Delta\hat{\sigma}_{\chi_0} = \frac{12\pi^2\alpha_S^2|R'_1|^2}{M_{\chi_0}^7}\delta\left(1 - \frac{M_{\chi_0}^2}{\hat{s}}\right), \quad \hat{\sigma}_{\chi_0} = \frac{12\pi^2\alpha_S^2|R'_1|^2}{M_{\chi_0}^7}\delta\left(1 - \frac{M_{\chi_0}^2}{\hat{s}}\right) \quad (6)$$

for χ_0 -productions and

$$\Delta\hat{\sigma}_{\chi_2} = -\frac{16\pi^2\alpha_S^2|R'_1|^2}{M_{\chi_2}^7}\delta\left(1 - \frac{M_{\chi_2}^2}{\hat{s}}\right), \quad \hat{\sigma}_{\chi_2} = \frac{16\pi^2\alpha_S^2|R'_1|^2}{M_{\chi_2}^7}\delta\left(1 - \frac{M_{\chi_2}^2}{\hat{s}}\right) \quad (7)$$

for χ_2 -productions. Here R'_1 is the derivative of the P-state wave function at the origin. The spin-dependent and spin-independent differential cross sections in terms of x_L ($= 2p_L/\sqrt{s}$) of produced χ_J ($J = 0, 2$) are

$$\frac{d\Delta\sigma_{\chi_J}}{dx_L} = \Delta C_{\chi_J} \frac{\pi^2 \alpha_S^2 |R'_1|^2}{M_{\chi_J}^7} \frac{\tau}{\sqrt{x_L^2 + 4\tau}} \delta G(x_a, Q^2) \delta G(x_b, Q^2), \quad (8)$$

$$\frac{d\sigma_{\chi_J}}{dx_L} = C_{\chi_J} \frac{\pi^2 \alpha_S^2 |R'_1|^2}{M_{\chi_J}^7} \frac{\tau}{\sqrt{x_L^2 + 4\tau}} G(x_a, Q^2) G(x_b, Q^2) \quad (9)$$

with x_a and x_b being the momentum fraction in a proton and given as

$$x_a = \frac{x_L + \sqrt{x_L^2 + 4\tau}}{2}, \quad x_b = \frac{-x_L + \sqrt{x_L^2 + 4\tau}}{2}, \quad \tau \equiv \frac{M_{\chi_{0,2}}^2}{s}, \quad (10)$$

where ΔC_{χ_J} and C_{χ_J} are 12 (−16) and 12 (16) for χ_0 (χ_2)-hadroproductions, respectively. By using these cross section formulas, the explicit form of two-spin asymmetries $A_{LL}^{\chi_{0(2)}}(pp)$ can be written as

$$A_{LL}^{\chi_{0(2)}}(pp) = (-) \frac{\delta G(\frac{x_L + \sqrt{x_L^2 + 4\tau}}{2}, Q^2)}{G(\frac{x_L + \sqrt{x_L^2 + 4\tau}}{2}, Q^2)} G(\frac{-x_L + \sqrt{x_L^2 + 4\tau}}{2}, Q^2). \quad (11)$$

It is remarkable that at low- p_T regions, $A_{LL}^{\chi_{0(2)}}(pp)$ which is originated mainly from gluon-gluon fusion of Figure 1, is directly proportional to polarized gluon distributions $\delta G(x)$. Therefore, $A_{LL}^{\chi_{0(2)}}(pp)$ sensitively depends on $\delta G(x)$. Namely, $A_{LL}^{\chi_{0(2)}}(pp)$ at low- p_T regions is a very good physical parameter to test the behavior of $\delta G(x)$. To derive the differential cross sections in Eqs.(8) and (9), the higher order QCD corrections, the quarkonium binding effect and the corrections due to relativistic effects of the quarkonium should be taken into account. However, these uncertainties are generally included in *so-called* “K-factor”, and they would cancel if those corrections contribute to the denominator and numerator of Eq.(11) alike.

As the PHENIX detector which is to work at RHIC is dedicated to have a special device for detecting leptons and γ 's[1], the observable signal for $\chi_{0,2}$ -productions is expected to be two-muon decays of 3S_1 coming from radiative decays $\chi_{0,2} \rightarrow ^3S_1 + \gamma$. Since the differential cross sections $d\Delta\sigma_{\chi_2}/dx_L$ and $d\sigma_{\chi_2}/dx_L$ are almost equal to $d\Delta\sigma_{\chi_0}/dx_L$ and $d\sigma_{\chi_0}/dx_L$, respectively, the χ_2 is easier to be detected because it has a larger branching ratio for the radiative decay than that of the χ_0 .

Using the polarized gluon distributions of four types given in Eqs.(1)–(4) and taking Q^2 as $M_{\chi_2}^2$, we estimate the $A_{LL}^{\chi_2}(pp)$ as a function of x_L for χ_2 -productions of not only the charmonium $\chi(c\bar{c})$ but also the bottomonium $\chi(b\bar{b})$ states at relevant

RHIC energies, $\sqrt{s} = 50, 200$ and 500GeV . Although the cross section of $\chi(b\bar{b})$ is smaller than that of $\chi(c\bar{c})$ about by 10^{-3} , $\chi(b\bar{b})$ has different values of x_a and x_b given in Eq.(10) from those of $\chi(c\bar{c})$ productions, and so we can obtain not only Q^2 -dependence but also x -dependence of the polarized gluon distributions by examining $\chi(b\bar{b})$ states together with $\chi(c\bar{c})$ states at the same time. We show the results of $A_{LL}^{\chi_2}$ calculated for (A) $c\bar{c}$ and (B) $b\bar{b}$ states at $\sqrt{s} = 50, 200$ and 500GeV in Figures 3, 4 and 5, respectively.

In the case of $\chi_2(c\bar{c})$, there is a tendency that the smaller \sqrt{s} (namely τ) is, the larger $A_{LL}^{\chi_2(c\bar{c})}$ become. In particular, for $\sqrt{s} = 50\text{GeV}$, it might be easy to examine not only the magnitude ΔG but also the behavior $\delta G(x)$ of the gluon polarization. In addition, $A_{LL}^{\chi_2(c\bar{c})}(pp)$ at $\sqrt{s} = 200$ and 500GeV is of use to know information on polarized gluon distributions, provided experiments are carried out with the high precision.

As for $\chi_2(b\bar{b})$ -productions, the dependence of $A_{LL}^{\chi_2(b\bar{b})}(pp)$ on \sqrt{s} has a similar tendency as that of $A_{LL}^{\chi_2(c\bar{c})}(pp)$. However, the behavior of $A_{LL}^{\chi_2(b\bar{b})}(pp)$ calculated at $\sqrt{s} = 50\text{GeV}$ by using our $\delta G(x)$ (type (a)) differs from that at the other \sqrt{s} . This is because for small x_L ($\lesssim 0.3$), x_a and x_b ($= x_a - x_L$) are taken to be about $0.2 - 0.4$ and hence $\delta G(x_a)/G(x_a)$ and $\delta G(x_b)/G(x_b)$ have around the minimal value as shown in Figure 2(C). Accordingly, Figure 3(B) indicates that it is rather difficult to distinguish our large $\delta G(x)$ (type (a)) from no polarization $\delta G(x) = 0$ (type (d)) in the region $x_L \lesssim 0.3$. At other \sqrt{s} , *i.e.* $\sqrt{s} = 200$ and 500GeV , since the difference of $A_{LL}^{\chi_2(b\bar{b})}$ for any type of polarized gluon distributions is relatively larger than that of $A_{LL}^{\chi_2(c\bar{c})}$, it might be useful to examine the magnitude and the x -dependence of δG , though the differential cross section of $\chi_2(b\bar{b})$ is smaller than that of $\chi_2(c\bar{c})$. Both $A_{LL}^{\chi_2(c\bar{c})}$ and $A_{LL}^{\chi_2(b\bar{b})}$ for the no gluon polarization (type (d)) are almost zero in Figures (4), (5) and (6).

We find that the low- p_T χ_2 -productions allow to give the clear test as a probe of the magnitude and the x -dependence of the gluon polarization because $A_{LL}^{\chi_2}$ for those reactions is directly proportional to $\delta G(x)/G(x)$. We have not taken up here the large- p_T χ_J -hadroproductions though they are also interesting for getting the informations on polarized gluon[18].

3 Summary

We have calculate the two-spin asymmetries for the low- p_T productions for χ_0 and χ_2 of the $c\bar{c}$ state and the $b\bar{b}$ state. The experimental data on charmonium and bottomonium χ_J -productions in polarized pp collisions at RHIC will be very helpful to put constraints on the magnitude and the behavior of the gluon polarization,

and improve our understanding of the proton structure. In particular, the low- p_T productions for χ_2 can give a very useful information on polarized gluon distributions because the two-spin asymmetry for these reactions is simply proportional to $\delta G(x)/G(x)$. We hope that our predictions will be tested in forthcoming RHIC spin experiments.

Acknowledgments

The authors are grateful to Profs. A. Masaike and S. Nurushev for useful comments on experimental issues at RHIC. We would like to acknowledge fruitful discussions on the subject of this work with Profs. G. Ramsey and D. Sivers.

References

- [1] RSC proposal, BNL, R 5, August (1992).
- [2] J. Ashman *et al.*, *Phys. Lett.* **B206**, 364 (1988); *Nucl. Phys.* **B328**, 1 (1989).
- [3] D. Adams *et al.*, *Phys. Lett.* **B329**, 399 (1994).
- [4] J. Ellis and R. L. Jaffe, *Phys. Rev.* **D9**, 1444 (1974); *Phys. Rev.* **D10**, 1669 (1974).
- [5] G. Altarelli and G. G. Ross, *Phys. Lett.* **B212**, 391 (1988); R. D. Carlitz, J. C. Collins and A. H. Mueller, *Phys. Lett.* **B214**, 229 (1988); A. V. Efremov and O. V. Teryaev, in *Proceedings of the International Hadron Symposium*, ed. Fischer *et al.*, (Czechoslovakian Academy of Science, Prague, 1989).
- [6] R. Baier and R. Rückl, *Z. Phys.* **C19**, 251 (1983); E. W. N. Glover, A. D. Martin and W. J. Stirling, *Z. Phys.* **C38**, 473 (1988).
- [7] G. Altarelli and W. J. Stirling, *Part. World* **1** 40, (1989); Z. Kunszt, *Phys. Lett.* **B218**, 243 (1989); J. Ellis, M. Karliner and C. T. Sachrajda, CERN-TH-5471/89; M. Glück, E. Reya and W. Vogelsang, *Phys. Rev.* **D45**, 2552 (1992).
- [8] H. Y. Cheng and S. N. Lai, *Phys. Rev.* **D41**, 91 (1990).
- [9] G. Ramsey and D. Sivers, *Phys. Rev.* **D43**, 2861 (1991).
- [10] T. Morii, S. Tanaka and T. Yamanishi, preprint KOBE-FHD-94-08 (1994), to be published in *Z. Phys.* **C**.
- [11] S. J. Brodsky, M. Burkardt and I. Schmidt, preprint 6087 (1994), to be published in *Nucl. Phys.* **B**.
- [12] T. Gehrmann and W. J. Stirling, Durham preprint DTP/94/38 (1994), to be published in *Z. Phys.* **C**.

- [13] D. L. Adams *et al.*, *Phys. Lett.* **B261**, 197 (1991).
- [14] D. L. Adams *et al.*, *Phys. Lett.* **B336**, 269 (1994).
- [15] W. Vogelsang and A. Weber, *Phys. Rev.* **D45**, 4069 (1992). T. Yamanishi, Doctoral thesis, Kobe University, March (1994).
- [16] J. L. Cortes and B. Pire, *Phys. Rev.* **D38**, 3586 (1988); M. A. Doncheski and R. W. Robinett, *Phys. Lett.* **B248**, 188 (1990).
- [17] R. Gastmans and T. T. Wu, *The Ubiquitous Photon Helicity Method for QED and QCD* (Oxford, 1990).
- [18] T. Morii, S. Tanaka and T. Yamanishi, preprint KOBE-FHD-95-02 (1995).

Figure captions

Fig. 1: A diagrams of the dominant process for low- p_T productions of $p p \rightarrow \chi_J X$ at lowest order QCD.

Fig. 2: The x -dependence of (A) $x\delta G(x, Q^2)$ at $Q^2 = 10\text{GeV}^2$, (B) $\delta G(x, Q^2)/G(x, Q^2)$ at $Q^2 = M_{\chi_2(c\bar{c})}^2$ and (C) $\delta G(x, Q^2)/G(x, Q^2)$ at $Q^2 = M_{\chi_2(b\bar{b})}^2$ for various types (a)–(d) given by Eqs.(1)–(4).

Fig. 3: Two-spin asymmetries (A) $A_{LL}^{\chi_2(c\bar{c})}(pp)$ and (B) $A_{LL}^{\chi_2(b\bar{b})}(pp)$ for $\sqrt{s} = 50\text{GeV}$, calculated with various types of $\delta G(x)$, as a function of longitudinal momentum fraction x_L of $\chi_2(c\bar{c})$ and $\chi_2(b\bar{b})$. The solid, dashed, small-dashed and dash-dotted lines indicate the results using types (a), (b), (c) and (d) in Eqs.(1), (2), (3) and (4), respectively.

Fig. 4: Two-spin asymmetries (A) $A_{LL}^{\chi_2(c\bar{c})}(pp)$ and (B) $A_{LL}^{\chi_2(b\bar{b})}(pp)$ for $\sqrt{s} = 200\text{GeV}$. Various lines represent the same as in Figure 3.

Fig. 5: Two-spin asymmetries (A) $A_{LL}^{\chi_2(c\bar{c})}(pp)$ and (B) $A_{LL}^{\chi_2(b\bar{b})}(pp)$ for $\sqrt{s} = 500\text{GeV}$. Various lines represent the same as in Figure 3.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9504251v1>

This figure "fig2-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9504251v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9504251v1>

This figure "fig2-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9504251v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9504251v1>

This figure "fig2-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9504251v1>